

Supplemental Table 1. Primer sequences for qPCR

Gene	Forward	Reverse	
<i>Wnt3</i>	GGGGCGTATTCAAGTAGCTG	GTAGGGACCTCCCATTGGAT	(1)
<i>Wnt6</i>	TGCCCCGAGGCGCAAGACTG	ATTGCAAACACGAAAGCTGTCTCTC	(2)
<i>Wnt9b</i>	AAGTACAGCACCAAGTTCCTCAGC	GAACAGCACAGGAGCCTGACAC	(2)
<i>Dll1</i>	CTGAGGTGTAAGATGGAAGCG	CAACTGTCCATAGTGCAATGG	(3)
<i>Dll4</i>	TCGTCGTCAGGGACAAGAATAGC	CTCGTCTGTTCCGCCAAATCTTACC	(3)
<i>Notch2</i>	TGCCTGTTTGACAACCTTTGAGT	GTGGTCTGCACAGTATTTGTCAT	(4)
<i>Hes1</i>	TGCCAGCTGATATAATGGAGAA	CCATGATAGGCTTTGATGACTTT	(4)
<i>Gapdh</i>	TTCCAGTATGACTCCACTCACGG	TGAAGACACCAGTAGACTCCACGAC	(5)

References

1. He W, et al. Wnt/beta-catenin signaling promotes renal interstitial fibrosis. *J Am Soc Nephrol.* 2009;20(4):765-76.
2. Davies PS, et al. Wnt-reporter expression pattern in the mouse intestine during homeostasis. *BMC Gastroenterology.* 2008;8(1):57.
3. Bohin N, et al. Rapid Crypt Cell Remodeling Regenerates the Intestinal Stem Cell Niche after Notch Inhibition. *Stem Cell Reports.* 2020;15(1):156-70.
4. Lim JS, et al. Intratumoural heterogeneity generated by Notch signalling promotes small-cell lung cancer. *Nature.* 2017;545(7654):360-4.
5. Ulatowski LM, et al. Strain-specific variants of the mouse *Cftr* promoter region reveal transcriptional regulatory elements. *Human Molecular Genetics.* 2004;13(17):1933-41.