

**Supplemental Table 1.** Baseline characteristics of those who returned for the follow-up OGTT (complete FUP) and those who did not (loos to FUP).

	High Incretin Effect		Moderate Incretin Effect		Low Incretin Effect	
	Complete FUP	Loss to FUP	Complete FUP	Loss to FUP	Complete FUP	Loss to FUP
<b>Baseline</b>	<b>Complete FUP</b>	<b>Loss to FUP</b>	<b>Complete FUP</b>	<b>Loss to FUP</b>	<b>Complete FUP</b>	<b>Loss to FUP</b>
<b>N</b>	<b>12</b>	<b>1</b>	<b>10</b>	<b>3</b>	<b>8</b>	<b>5</b>
<b>Age (years)</b>	15.7 (14.5, 17.1)	18 (18, 18)	16.2 (15, 17)	13(12, 14)	16.3 (14.0, 19.2)	15 (12, 16)
<b>BMI (kg/m<sup>2</sup>)</b>	32.5 (29.8, 40.3)	44.9 (44.9, 44.9)	37.2 (35.7, 40.8)	41.0 (35.3, 42.6)	41 (34.0, 45.3)	39.8 (37, 41.4)
<b>NGT/preIGT/IGT/T2D n (%)</b>	0/9/3/0 (0/67/33/0)	0/0/1/0 (0/0/100/0)	0/5/5/0 (0/50/50/0)	0/2/1/0 (0/70/30/0)	0/4/4/0 (0/38/62/0)	0/1/4/0 (0/20/80/2)
<b>Sex (F/M/Non-binary) (%)</b>	9/3/0 (69/21/0)	1/0/0 (100/0/0)	3/7/0 (31/69/0)	2/1/0 (67/33)	4/4/0 (50/50/0)	3/2/0 (60/40/0)
<b>Ethnicity NHW/NHB/H n (%)</b>	1/5/6 (8/42/50)	0/1/0 (0/100/0)	1/5/4 (10/50/40)	0/1/2 (0/33/67)	3/1/4 (37/13/50)	4/0/1 (80/0/20)

NGT, normal glucose tolerance; pre-IGT, pre-impaired glucose tolerance; IGT, impaired glucose tolerance; T2D, type 2 diabetes; BMI, body mass index; NHW, non-Hispanic White; NHB, non-Hispanic Black; H, Hispanic; FUP, follow-up.

