


Sparsentan improves glomerular hemodynamics, cell functions and tissue repair in a mouse model of FSGS

Georgina Gyarmati, ... , Patricia W. Bedard, Janos Peti-Peterdi

JCI Insight. 2024. <https://doi.org/10.1172/jci.insight.177775>.

Research In-Press Preview Nephrology

Graphical abstract



Find the latest version:

<https://jci.me/177775/pdf>



Sparsentan improves glomerular hemodynamics, cell functions and tissue repair in a mouse model of FSGS

Georgina Gyarmati¹, Urvi Nikhil Shroff¹, Audrey Izuhara¹, Sachin Deepak¹, Radko Komers², Patricia W. Bedard², Janos Peti-Peterdi¹

¹Departments of Physiology and Neuroscience, and Medicine, Zilkha Neurogenetic Institute, University of Southern California, Los Angeles, California, USA

²Travere Therapeutics, San Diego, California, USA

Corresponding author: Janos Peti-Peterdi, Zilkha Neurogenetic Institute, Room ZNI313, 1501 San Pablo Street, Los Angeles, CA 90033. Phone: (323)442-4337. Email: petipete@usc.edu

Conflict of interest statement:

This study was funded by Travere Therapeutics.

ABSTRACT

Dual endothelin-1 (ET-1) and angiotensin II (AngII) receptor antagonism with sparsentan has strong antiproteinuric actions via multiple potential mechanisms that are more pronounced, or additive compared to current standard of care using angiotensin receptor blockers (ARB). Considering the many actions of ET-1 and AngII on multiple cell types, this study aimed to determine glomeruloprotective mechanisms of sparsentan compared to the ARB losartan by direct visualization of its effects in the intact kidney in focal segmental glomerulosclerosis (FSGS) using intravital multiphoton microscopy. In both healthy and FSGS models, sparsentan treatment increased afferent/efferent arteriole diameters, increased or preserved blood flow and single nephron glomerular filtration rate, attenuated acute ET-1+AngII-induced increases in podocyte calcium, reduced proteinuria, preserved podocyte number, increased both endothelial and renin lineage cells and clones in vasculature, glomeruli and tubules, restored glomerular endothelial glycocalyx, attenuated mitochondrial stress and immune cell homing. These effects were either not observed or of smaller magnitude with losartan. The pleiotropic nephroprotective effects of sparsentan included improved hemodynamics, podocyte and endothelial cell functions, and tissue repair. Compared to losartan, sparsentan was more effective in the sustained preservation of kidney structure and function, which underscores the importance of the ET-1 component in FSGS pathogenesis and therapy.

INTRODUCTION

Focal segmental glomerulosclerosis (FSGS) refers to the typical focal and segmental renal histopathology pattern of glomerular scarring that represents a variety of kidney diseases that may result in progressive chronic kidney disease (CKD) and end stage kidney disease (ESKD). FSGS accounts for about 5% of the ~750,000 US adults with ESKD (1). Most primary, secondary and genetic causes of FSGS are associated with the injury and loss of podocytes, a special cell type of the glomerular filtration barrier (GFB). In cases of genetic FSGS the lesions develop due to defects in various proteins of the podocyte actin-cytoskeleton and slit diaphragm complex (2, 3). Recent mechanistic insights based on genetic mutations, multiple animal models and human clinical studies highlighted the pathogenic role of at least one such protein called TRPC6 (Transient Receptor Potential cation channel, subfamily C, member 6), that mediates podocyte Ca^{2+} influx (4-7). TRPC6 gain-of-function mutations augment AngII-dependent increases in podocyte calcium, causing dysfunction and loss of podocytes and hereditary FSGS (7, 8). A mouse model with ~4-fold overexpression of wild-type TRPC6 only in podocytes develops human FSGS-like kidney disease, which has been a useful tool for research (5). In both humans and animal models of glomerulosclerosis, albuminuria is the first sign of glomerular dysfunction, which is often later associated with renal function decline and tissue sclerosis (3). Many patients with FSGS have a progressive course of CKD and ultimately require renal replacement therapy, until new and highly efficacious therapies are developed that can stop or reverse kidney function decline.

Sparsentan is a first-in-class, orally active, single-molecule dual selective antagonist of the angiotensin II subtype 1 receptor (AT_1R) and the Endothelin A receptor ($\text{ET}_\text{A}\text{R}$) (9, 10). Results of recent clinical studies demonstrated strong antiproteinuric actions of sparsentan versus the ARB irbesartan in patients with FSGS (11-14) and also IgA nephropathy (15), suggesting

nephroprotective effects of the drug. Considering the large spectrum of renal actions of AngII and ET-1 including vasoconstriction of the afferent (AA) and efferent arterioles (EA) and mesangial contraction that reduce blood flow and glomerular filtration rate, and their upregulation in kidney injury including FSGS (13), dual inhibition of their actions, as occurs when using sparsentan, is expected to exert multiple actions in many renal cell types and consequently result in a variety of renoprotective mechanisms. Based on recently published and emerging evidence in different models of glomerular disease, sparsentan may preserve glomerular filtration rate and GFB structure and function, inhibit renal cell growth and extracellular matrix production together with anti-sclerotic, anti-fibrotic and anti-inflammatory mechanisms, and may augment endogenous tissue repair (13, 16). However, the clear nature of these mechanisms and consequently the primary modes of action of sparsentan, in particular in proteinuric kidney diseases are not yet completely understood. One major bottleneck of progress in our improved mechanistic understanding of FSGS pathophysiology and therapeutic mechanisms has been the inability to study the structurally and functionally highly complex kidney tissue in its native environment. To advance the field, the present study applied a direct intravital visual approach and comprehensive analysis to study the complexity of sparsentan actions on hemodynamics, cell biology functions, and tissue remodeling.

Recent advances in high-power intravital imaging using multiphoton microscopy (MPM) have made it possible to directly visualize over time the development and progression of glomerular disease processes in the intact mouse kidney in vivo in unprecedented detail (17-23). In addition, transgenic mouse models are available that allow renal cell-specific expression of fluorescent reporters (e.g. the multi-color Confetti construct)(20, 24-26) for genetic cell fate tracking and to study single cell-based changes in functional (e.g. intracellular calcium)(18, 25, 26) and ultrastructural parameters of the GFB, renal tubules and microvessels with subcellular

resolution in combination with in vivo MPM imaging (22, 25-28). The unique ability of this approach to be able to measure cell calcium in podocytes, afferent (AA) and efferent arteriole (EA) vascular smooth muscle cells (VSMC)(18, 25, 26), albumin leakage through the GFB(21, 22), AA/EA/glomerular capillary diameter (19, 29), glomerular endothelial surface layer (19, 21, 23, 30), and immune cell homing (21, 31, 32) has been demonstrated previously.

The present study used intravital MPM imaging of genetically engineered mouse models with fluorescent lineage tags and calcium reporters in various kidney cell types (podocytes, endothelial and vascular progenitor and smooth muscle cells, and tubular epithelia) to quantitatively visualize the mode of action and compare the effects of sparsentan and the ARB losartan on glomerular hemodynamics, GFB function and kidney tissue remodeling under physiologic conditions and in FSGS.

RESULTS

Effects of sparsentan on glomerular hemodynamics under physiological conditions

We first tested the effects of sparsentan as compared to losartan on glomerular hemodynamics in the normal healthy kidney (Figure 1A-I). Ren1d-GCaMP5/tdTomato mice (6-8 weeks old) that express the calcium reporter GCaMP5 and the calcium non-sensitive tdTomato in cells of the renin lineage (including AA/EA VSMC and intraglomerular mesangial cells, Figure 1B) were treated chronically with either no drug control, sparsentan (120 mg/kg body wt), or losartan (10 mg/kg body wt) daily for six weeks and prepared for intravital MPM imaging of numerous structural and functional parameters of glomerular hemodynamics at the single nephron level. Sparsentan (6-week treatment) markedly increased AA (20.23 ± 0.76 vs 14.94 ± 0.43 μm in control, Figure 1C) and EA diameters (12.30 ± 0.98 vs 9.59 ± 0.64 μm in control, Figure 1D) and

single nephron glomerular filtration rate (SNGFR) (6.36 ± 0.50 vs 4.57 ± 0.38 nl/min in control, Figure 1H). No changes were observed in other hemodynamic parameters including glomerular diameter or tuft area, glomerular capillary diameter and red blood cell (RBC) velocity (Figure 1E-I). Losartan did not cause changes in any of these glomerular hemodynamic parameters (Figure 1C-I).

In addition, we tested the ability of sparsentan to prevent glomerular hemodynamic alterations caused by acute agonist-induced vasoconstriction. Endothelin (ET-1, 50 ng/kg) with or without AngII (400 ng/kg) was injected in bolus into the cannulated carotid artery of the same Ren1d-GCaMP5/tdTomato mice (Figure 2A) that were pre-treated with either no drug control, sparsentan (120 mg/kg), or losartan (10 mg/kg) for six weeks. In control mice, combined injection of ET-1+AngII caused strong vasoconstriction of the AA ($51.93 \pm 5.62\%$ of baseline diameter) which was also confirmed by the substantial elevations in AA VSMC calcium (5.90 ± 0.59 fold of baseline; Figure 2A-D, and Supplementary Video 1). The glomerular diameter ($91.61 \pm 2.80\%$ of baseline diameter) and tuft area ($92.79 \pm 0.99\%$ of baseline diameter) reduced similarly (Figure 2E). While losartan inhibited only the elevations in dual agonist-induced AA VSMC calcium (3.27 ± 0.21 fold of baseline, Figure 2B-C) but not the other hemodynamic parameters (Figure 2D-F), sparsentan almost completely abolished dual agonist-induced AA VSMC calcium elevations (1.93 ± 0.19 fold of baseline) and AA vasoconstriction ($86.41 \pm 4.44\%$ of baseline diameter, Figure 2B-D, and Supplementary Video 1). These effects of sparsentan were more pronounced than those of losartan (Figure 2B-F). While sparsentan partially attenuated the agonist effects on glomerular diameter and tuft area, these effects did not reach statistical significance (Figure 2E-F). ET-1 injection alone produced similarly strong AA/glomerular contractions (Figure 2C-F). While losartan had no effect on any of the glomerular hemodynamic alterations (Figure 2C-F), treatment

with sparsentan almost completely abolished ET-1-induced elevations in AA VSMC calcium (1.41 ± 0.17 fold of baseline compared 3.36 ± 0.27 in control), AA vasoconstriction ($93.26 \pm 2.85\%$ of baseline diameter compared to $49.35 \pm 4.62\%$ in control), and the reduction in glomerular tuft area ($97.30 \pm 1.14\%$ of baseline compared to $88.31 \pm 2.58\%$ in control, Figure 2C-F). In addition to the effects on AA VSMCs, sparsentan almost completely abolished the ET-1 injection induced elevations in intracellular calcium also in intraglomerular mesangial cells (1.52 ± 0.07 fold of baseline compared 4.52 ± 0.46 in control) and in cells of the distal convoluted tubule (1.17 ± 0.10 fold of baseline compared 2.71 ± 0.32 in control) (Supplementary Video 2). The distal tubule effect of ET-1 was likely mediated via the actions of filtered ET-1 in the tubular fluid acting on ET-1 receptors localized at the apical cell membrane, based on the delayed nature of this effect appearing 30-40 s after injection and with a calcium wave propagation pattern from the apical towards basal cell regions (Supplementary Video 2).

Effects of sparsentan on kidney tissue remodeling under physiological conditions

To test the effects of sparsentan compared to losartan on endogenous kidney tissue remodeling in the normal healthy kidney, Ren1d-Confetti and Cdh5-Confetti mice (6-8 weeks old) were used that express the multi-color Confetti reporter (membrane CFP (blue), nuclear GFP (green), cytosolic YFP (yellow) and RFP (red)) in cells of the renin lineage or endothelium, respectively. Mice were treated with either no drug control, sparsentan (120 mg/kg), or losartan (10 mg/kg) daily for two weeks and then prepared for histological analysis of fixed kidney sections. In Ren1d-Confetti mice, sparsentan treatment resulted in a substantial increase in the number of Confetti+ cells (18.19 ± 2.23 compared to 5.75 ± 0.28 in control), the number of identical Confetti color cell groups (clones) (2.50 ± 0.16 compared to 0.30 ± 0.07 in control), and the number of

individual Confetti⁺ cells per clone (9.13 ± 0.79 compared to 4.56 ± 0.11 in control) in the glomerular tuft, at the glomerular vascular pole and terminal AA segment (Figure 3A-D). Sparsentan had a similar effect on clonal remodeling of the glomerular and vascular endothelium (Cdh5-Confetti mice, Figure 3F-J). In summary, sparsentan increased the frequency of larger multi-cell clones in both the renin and endothelial lineage (Figure 3E, J). Losartan had a similar effect on both the Ren1d and Cdh5-Confetti cell populations, but with a reduced magnitude compared to sparsentan (Figure 3A-J). In addition, various renal cortical and medullary tubule segments including cells of the proximal tubule, the distal convoluted tubule and the collecting duct also showed active cellular (clonal) remodeling in response to sparsentan based on the increased frequency of larger multi-cell clones, and with a modest response to losartan (Figure 3K-L).

To confirm that sparsentan and losartan administration via custom-made mouse chow in the present study resulted in therapeutically relevant and effective plasma levels, two blood samples (early morning and late afternoon consistent with the 12 hours dark/light cycle) were collected from all animals at the end of the six weeks treatment for the measurement of plasma drug levels. Sparsentan and losartan were detectable in the plasma in all mice and time points (data not shown), and there was no difference in systolic blood pressure between the three treatment groups in physiological conditions measured by tail-cuff (111.2 ± 5.7 mmHg in control, 106.4 ± 2.3 mmHg in sparsentan ($p=0.16$ vs control) and 112.2 ± 5.1 mmHg in losartan ($p=0.92$ vs control) treatment).

Effects of sparsentan on glomerular hemodynamics and GFB function in FSGS

We next evaluated the effects of sparsentan as compared to losartan on glomerular hemodynamics under disease conditions using a mouse model of FSGS. Pod-GCaMP5/tdTomato TRPC6 TG mice were developed that express the calcium reporter GCaMP5 and the calcium non-sensitive tdTomato only in podocytes, and also feature ~4-fold overexpression of TRPC6 in podocytes. These mice (1.5 years old) that are at an advanced age when substantial pathology is present, as reported previously (5), were treated chronically with either no drug control, sparsentan (120 mg/kg), or losartan (10 mg/kg) daily for six weeks and prepared for intravital MPM imaging of the same glomerular hemodynamic parameters as before in the physiological model. In contrast to the healthy physiological state and similarly to another pre-clinical FSGS model using Pod-GCaMP3 mice in our earlier study (18), in no drug control, Pod-GCaMP5/tdTomato TRPC6 TG mice had the visual signs of ongoing FSGS pathology including segmental elevations in podocyte calcium, the development of multiple parietal podocytes and adhesions between parietal Bowman's capsule and glomerular capillary segments, and albumin leakage through the GFB (Figure 4A-J). Sparsentan treatment markedly improved several parameters of glomerular hemodynamics and GFB function compared to control (Figure 4B-J), including reductions in podocyte calcium (GCaMP5/tdTomato fluorescence ratio 0.16 ± 0.02 vs 1.1 ± 0.19 in control, Figure 4B), increased AA (17.92 ± 0.47 vs 11.65 ± 0.57 μm in control) and EA diameters (10.39 ± 0.40 vs 7.53 ± 0.69 μm in control, Figure 4C-D), SNGFR (8.22 ± 0.48 vs 3.11 ± 0.29 nl/min in control, Figure 4G), and glomerular capillary blood flow (RBC velocity 2.29 ± 0.19 vs 0.88 ± 0.15 $\mu\text{m}/\text{ms}$ in control, Figure 4H). In contrast to the low glomerular albumin permeability in control healthy conditions (albumin GSC below 0.015) as established in previous in vivo MPM imaging studies (33, 34), the presently applied FSGS model featured glomerular albumin leakage (Figure 4I). Importantly, albumin leakage through the GFB (albumin GSC 0.10 ± 0.02 vs 0.22 ± 0.03 in control, Figure 4I),

and the level of albuminuria (urinary albumin/creatinine ratio, ACR normalized to baseline 0.71 ± 0.04 vs 2.52 ± 0.76 in control, Figure 4J) also improved markedly in response to sparsentan treatment. Of all the parameters measured, losartan improved only podocyte calcium (Figure 4B) and glomerular capillary blood flow (Figure 4H), but with a lower magnitude compared to sparsentan.

At the end of the chronic treatment for six weeks in 6-12 months old TRPC6 TG (FSGS) mice, systolic blood pressure was measured at the end of the resting (non-feeding, lowest expected plasma drug levels) phase of their circadian cycle. Direct measurements by pressure transducer via the cannulated carotid artery in anesthetized animals showed equal reduction in blood pressure by sparsentan (113.2 ± 0.4 mmHg) and losartan (108.5 ± 2.0 mmHg) compared to control (129.2 ± 1.4 mmHg) (Figure 4K). To further confirm that the extent of AT₁R blockade was equal between sparsentan and losartan treatment, systolic blood pressure responses to acute AngII injection were measured via pressure transducer. In control mice, acute bolus injections of AngII (400 ng/kg) caused marked elevations in systolic blood pressure (Δ BP 43.1 ± 1.6 mmHg), which were substantially and equally lowered in mice chronically treated with sparsentan (30.6 ± 2.5 mmHg) or losartan (30.6 ± 3.1 mmHg) (Figure 4L).

Additional experiments were performed in FSGS mice to test the ability of sparsentan to prevent the alterations in glomerular hemodynamics and GFB function caused by acute agonist-induced vasoconstriction, the same way it was tested in the control healthy kidney (Figure 2). ET-1 (50 ng/kg) and AngII (400 ng/kg) were mixed together and injected in bolus into the cannulated carotid artery of the same Pod-GCaMP5/tdTomato TRPC6 TG mice that were pre-treated with either no drug control, sparsentan (120 mg/kg), or losartan (10 mg/kg) for six weeks. In control mice, ET-1+AngII injections produced major elevations in podocyte calcium (3.88 ± 0.66 fold of

baseline), strong vasoconstriction of the AA ($66.31 \pm 3.4\%$ of baseline diameter), reduction in glomerular diameter ($93.99 \pm 0.88\%$ of baseline) and tuft area ($90.65 \pm 1.85\%$ of baseline) (Figure 5A-E). While losartan had no effect on most of these parameters except on reducing agonist-induced podocyte calcium elevations (Figure 5B) and glomerular tuft area (Figure 5E), treatment with sparsentan almost completely abolished ET-1+AngII-induced podocyte calcium elevations (1.02 ± 0.05 fold of baseline), AA vasoconstriction ($88.67 \pm 3.85\%$ of baseline diameter), reductions in glomerular diameter ($98.14 \pm 0.48\%$ of baseline) and glomerular tuft area ($97.00 \pm 0.62\%$ of baseline, Figure 5A-E).

Effects of sparsentan on kidney tissue remodeling in FSGS

Traditional histology-based phenotyping of podocyte number, glomerulosclerosis and tissue fibrosis was performed from harvested Pod-GCaMP5/tdTomato TRPC6 TG mouse kidneys in the three treatment groups. Compared to the normally high podocyte number in healthy, non-FSGS mice as established in our recent study using the same protocol (31), podocyte number was low in untreated TRPC6 TG FSGS mice (Figure 6A). Importantly, the results confirmed the therapeutic benefit of sparsentan on preserving p57⁺ (12.46 ± 0.57 vs 4.37 ± 0.56 in control, Figure 6A) and WT1⁺ podocyte number (19.46 ± 0.35 vs 7.02 ± 0.19 in control, Figure 6B) and reducing glomerulosclerosis (47.17 ± 3.20 vs 101.70 ± 7.15 in control) and tissue fibrosis (36.30 ± 1.77 vs 86.23 ± 4.42 in control, Figure 6C). Although losartan also improved these parameters, it was less effective in preserving podocyte number compared to sparsentan (Figure 6A-C).

To test the effects of sparsentan compared to losartan on endogenous kidney tissue remodeling in FSGS as described in the physiological condition (Figure 3), Ren1d-Confetti and Cdh5-Confetti mice were crossed with Pod-TRPC6 TG mice to track the fate of cells of the renin

lineage or endothelium, respectively. Pod-TRPC6TG/Ren1d-Confetti and Pod-TRPC6TG/Cdh5-Confetti mice (6 months old) were treated chronically with either no drug control, sparsentan (120 mg/kg), or losartan (10 mg/kg) daily for six weeks and then prepared for histological analysis. Sparsentan treatment resulted in substantial increases in the number of Ren1d-Confetti+ cells (29.38 ± 1.22 compared to 14.75 ± 1.74 in control), the number of identical Confetti color cell groups (clones) (2.56 ± 0.30 compared to 0.25 ± 0.11 in control), and the number of individual Confetti+ cells per clone (10.39 ± 1.63 compared to 1.38 ± 0.32 in control) in glomeruli including the vascular pole and terminal AA segment (Figure 6D). Sparsentan had a similar effect on clonal remodeling of the glomerular and vascular endothelium in FSGS (Figure 6E). In sum, sparsentan increased the frequency of larger multi-cell clones in both the renin and endothelial lineage in FSGS (Figure 6F, G). Entirely clonal (unicolor) glomeruli and arterioles were often observed in response to sparsentan but not losartan treatment (Figure 6D-E). Losartan had a similar effect on both the Ren1d and Cdh5-Confetti cell populations, but with a reduced magnitude compared to sparsentan (Figure 6D-G).

Effects of sparsentan on glomerular cell metabolism, endothelial surface layer and immune cell homing in FSGS

To determine specific glomerular functional effects of sparsentan at the cell and molecular level, chronically treated Pod-TRPC6 TG mice as described above were injected with either MitoTracker Red or FITC-WGA and anti-CD44-Alexa Fluor 488 antibodies to quantitatively visualize alterations in cell metabolism, endothelial surface layer (glycocalyx) density, and glomerular immune cell homing, respectively. Sparsentan treatment resulted in substantial reduction of the mitochondrial membrane potential (an index of oxidative stress) in all

glomerular cell types including podocytes (815.3 ± 52.10 AU compared to 1258 ± 88.96 AU in control, Figure 7A). Sparsentan also improved the glomerular endothelial surface layer compared to the heterogeneous, segmentally high or undetectable levels observed in control (as a sign of endothelial dysfunction (31)) (0.72 ± 0.03 μm compared to 1.14 ± 0.08 μm in control, Figure 7B). In addition, numerous immune cells were observed homed to the glomerular capillary lumen in control (as a sign of local inflammation), that were eliminated by sparsentan (4.25 ± 0.70 compared to 9.94 ± 0.84 in control, Figure 7B). Losartan had a similar effect on most but not all these glomerular cell functions compared to sparsentan (Figure 7A-B).

DISCUSSION

The present study applied an intravital imaging approach using MPM of the local kidney tissue microenvironment at the single nephron level to directly and quantitatively visualize the effects of sparsentan in the control healthy kidney and in a disease model of FSGS. The key findings were the broad spectrum and positive classic hemodynamic and novel tissue regenerative, cell metabolism and endothelial protective and anti-inflammatory effects of sparsentan, and in a number of parameters their superiority in comparison to the ARB losartan. This study represents a major advance towards our improved mechanistic understanding of the specific mode of actions of sparsentan. It also demonstrated the importance of the additive effects or interplay of ET-1 and AngII in the maintenance of renal hemodynamic and glomerular cell biological functions and tissue remodeling in physiological conditions and in the pathogenesis and treatment of FSGS.

In addition to a physiological model, the present study used a disease model, aged mice with podocyte-specific overexpression of wild type TRPC6 which was shown to result in human

FSGS-like disease (5). The human clinical relevance due to pathological features and genetic cause of FSGS due to TRPC6 gain-of-function makes this mouse model very useful for the pre-clinical testing of sparsentan. The combination of TRPC6 TG with cell-specific expression of fluorescent calcium and cell lineage reporters in podocytes, vascular smooth muscle, endothelial and mesenchymal progenitor cells was highly valuable in the present study for the direct visualization of the acute or chronic effects of sparsentan or losartan and the actions of ET-1 with or without AngII on these renal cell types. Several molecular and cellular mechanistic details of the mode of action of sparsentan were uncovered in both control healthy and FSGS disease conditions. The results from sparsentan and losartan treatments in physiological models suggest that AngII and ET-1 are active, intrinsic players and important determinants of glomerular hemodynamics and physiological tissue maintenance.

In patients with FSGS, sparsentan lowered proteinuria at 108 weeks of treatment versus irbesartan (14), however, the mechanistic basis for the antiproteinuric effect of sparsentan versus ARB was not addressed in that study and remains to be elucidated in more detail. As the present MPM imaging approach directly and visually demonstrated (Figures 4-7), a combination of several factors including improved glomerular hemodynamics, cell biological functional, and tissue remodeling mechanisms were likely involved in the protective antiproteinuric effects of sparsentan. Importantly, chronic treatment with sparsentan preserved glomerular hemodynamics and GFB functions in FSGS, as well as ameliorated the strong effects of acute vasoconstrictor challenges with the two mechanistically and therapeutically relevant agonists (ET-1 and AngII). In contrast to losartan, sparsentan-treated animals demonstrated greater AA/EA diameters and higher SNGFR in physiological as well as FSGS disease conditions (Figures 1 and 4) suggesting that ET-1 is an important contributor to the regulation of glomerular hemodynamics in the healthy

kidney and a player in FSGS pathogenesis. Because sparsentan dilated both the AA and EA and increased glomerular capillary RBC velocity (blood flow) considerably (Figure 4H), the increased SNGFR was likely due to an increased filtration coefficient (GFB surface area and permeability) rather than increased glomerular capillary hydrostatic pressure, although this was not measured directly in our study. In addition, increased glomerular blood flow is known to attenuate the rise in oncotic pressure along the glomerular capillary and thereby enhances effective net filtration pressure and SNGFR. The glomerular endothelial remodeling effects (Figures 3 and 6) and the observed trend to increased glomerular tuft and diameter (Figure 4E-F) with sparsentan treatment are consistent with the above notions. In addition, the reduction in podocyte intracellular calcium levels in response to sparsentan (Figure 4B) is an indication of reduced rather than increased mechanical strain on the GFB, assuming that sparsentan does not directly/indirectly interfere with podocyte calcium handling. The more pronounced effects of sparsentan to abolish acute agonist-induced alterations in glomerular hemodynamics (Figure 2) are consistent with the additive protective effect of ET-1 receptor blockade on top of ARB. Several protective effects of losartan confirmed the well-established pathogenic role of AngII in CKD and FSGS, including the reduced AA VSMC (Figure 2C) and podocyte intracellular calcium levels (Figure 4B), improved glomerular capillary blood flow (Figure 4H), increased podocyte number and reduced glomerulosclerosis and tissue fibrosis (Figure 6A-C). The similar effects of sparsentan and losartan on glomerulosclerosis and tissue fibrosis (Figure 6C) despite the clear benefit of sparsentan vs losartan treatment on other parameters such as podocyte number (Figure 6A-B) suggest that 6 weeks of treatment may not be sufficiently long or the disease stage was too advanced (mice were 1.5 years old) to translate into improved histological readouts in a slowly developing FSGS model such as the applied TRPC6 transgenic mouse model. Although it remains to be determined if the

hemodynamic data measured at the single nephron level translate to global GFR changes, the results strongly suggest that improving renal hemodynamics is one of important modes of action of the therapeutic benefit of dual ET-1 and AngII antagonism with sparsentan. In addition, we cannot exclude that glomerular hemodynamic differences between the sparsentan- and losartan-treated animals are at least in part attributable to better preservation of glomerular structural integrity in the sparsentan treatment group. Interestingly, the present study uncovered the direct effects of ET-1 and sparsentan not only on the AA and EA, but also within the glomerulus on intraglomerular mesangial cells and in numerous renal tubular segments (Figures 1-2, Supplement Videos 1-2). Decreased intracellular calcium in mesangial cells of sparsentan-treated animals is consistent with decreased activation and proliferation seen in the gddY and EIC mouse models of IgA nephropathy (35, 36). These findings are consistent with the expression of the ET_AR and AT₁R in these vascular, glomerular, and tubular cell types (37, 38).

Although very few glomerular hemodynamic and GFB parameters were altered by losartan treatment, AT₁R blockade by losartan inhibited ET-1+AngII-induced elevations in AA VSMC (in physiological model) and podocyte (in FSGS model) calcium (Figures 2C and 4B) and improved glomerular capillary blood flow in the FSGS model (Figure 4H). The observed losartan effect on reducing intracellular calcium but not enough to cause vasodilation (without changing vascular diameter, Figure 2C-D) is consistent with the much higher sensitivity of the presently applied novel MPM calcium imaging approach compared to the classic dimensional readouts (vascular diameter) of AA contractility. The lack of AA vasodilation in response to losartan (Figure 2D) may be explained by the preferential effect of AngII on the EA rather than the AA. In these acute vasoconstrictor injection experiments the potential changes in EA diameter could not be measured simultaneously due to technical limitations. The increased blood flow without changes in AA/EA

diameter in response to losartan (Figure 4C-D, H) may be explained by preferential endothelial actions that are supported by the presently observed endothelial protective effects (Figure 7A-B). These results are consistent with the measured plasma losartan drug levels which were found to be detectable and in the therapeutic range. However, the more robust and broad effects of sparsentan on many glomerular hemodynamic and GFB parameters (Figures 1-2,4,6) suggest the primary importance of interplay of ET-1 and AngII signaling in both the physiological maintenance and disease alterations in glomerular hemodynamics and GFB functions. It needs to be emphasized that the extent and daily duration of AT₁R blockade were equal with both sparsentan and losartan treatment. This was indicated by blood pressure data that were similar in the physiological model and equally reduced in FSGS conditions by both sparsentan and losartan compared to control (Figure 4K). Importantly, the acute AngII-induced blood pressure elevations were also equally lowered by both sparsentan and losartan compared to control (Figure 4L). These results further underscore the importance and therapeutic benefit of the ET-1 component of sparsentan and its interplay with AngII signaling in FSGS pathogenesis, independently of blood pressure changes.

The substantial augmentation of protective endogenous tissue remodeling by sparsentan that involves the cells of the renin and endothelial cell lineages (Figures 3 and 6) is an unexpected finding. The renin cell lineage is known to have the ability to function as progenitor cells for parietal epithelial cells and podocytes in glomerular structural and functional repair (39, 40). The results suggest that ET-1 and AngII signaling are major regulators of promoting differentiation rather than self-renewal of renin and endothelial progenitor cells. Based on the superior effects of dual ET-1+Ang II receptor blockade with sparsentan versus inhibition of only AngII receptors with losartan (Figures 3 and 6), ET-1 rather than AngII signaling may be of primary importance in this function. Sparsentan treatment comparably augmented progenitor cell-mediated tissue

remodeling in healthy mice (Figure 3) and regeneration in FSGS (Figure 6), suggesting the preservation of its “physiological” tissue remodeling capacity in the disease setting. Importantly, this effect was more pronounced than those of losartan even if renin-angiotensin system inhibition is known to potently increase podocyte derivation from cells of the renin lineage (39). The effects of sparsentan to increase the number of Ren1d-Confetti+ clones and the number of cells within one clone (Figure 3C-E) are consistent with its direct effects on progenitor cells and their proliferation. The use of two different podocyte markers (p57⁺ and WT1⁺) found preservation of podocyte number by sparsentan and less potently by losartan (Figure 6A-B) within only a relatively short (6 weeks) treatment compared to the 1.5 years of age and ongoing FSGS pathology development. These results suggest that the recruitment of new podocytes from renin/PEC progenitors (41-43) rather than reduced cell death was involved in this phenomenon. RAS inhibition and low dietary salt intake that are both associated with increases in the number of renin lineage cells and are known to reduce albuminuria in CKD via glomerular structural improvements independent of blood pressure changes (44). In addition to the renin lineage, the recent development of the Cdh5-Confetti model for genetic fate tracking of endothelial cells at the single-cell level (30, 45) enabled testing of the effect of sparsentan on endothelial precursor cells in the present study. Similarly to renin cells, sparsentan increased the number of Cdh5-Confetti+ clones, the number of cells within one clone, and the frequency of clones with high cell number (Figure 3F-J). These results are consistent with the direct effects of sparsentan on endothelial precursor cells and their proliferation. Again, the superior effects of dual ET-1+Ang II receptor blockade compared to the ARB losartan on renal vascular, glomerular, and tubular remodeling and regeneration is an exciting finding. These results strongly suggest that in addition to its effects on

renal hemodynamics, kidney tissue regenerative mode of actions are involved and are at least as important in the superior therapeutic benefit by sparsentan treatment.

Glomerular disease-relevant actions of both ET-1 and AngII are known to include not only podocyte calcium signaling (Figures 4B and 5B) but also the generation of reactive oxygen species, oxidative stress, inflammation, and degradation of the glomerular endothelial surface layer in models of FSGS (46-50). Therefore, additional *in vivo* MPM imaging studies were performed to test the effects of sparsentan on these glomerular cell and molecular targets. Similarly to the beneficial effects seen earlier in this study, the results confirmed the therapeutic benefit of sparsentan (in most cases with a trend to be superior to losartan) on normalizing the excessive mitochondrial metabolism (oxidative stress) in podocytes and other glomerular cell types (Figure 7A), the endothelial surface layer in glomerular endothelial cells (Figure 7B) and reducing the glomerular homing of immune cells (Figure 7B). The endothelial glycocalyx protective effects of sparsentan are consistent with recent reports on the similar effects of sparsentan in a different animal disease model or those of ET_A receptor inhibition (35, 51). All parameters measured in the present study were tested for the potential impact of sex as a biological variable, and none of the results found difference between males and females (Figures 1-7).

In summary, the present study provides direct *in vivo* data that point to multiple layers of protective effects of sparsentan in a model of FSGS demonstrating improvements in glomerular hemodynamics, restoration of the glomerular endothelial surface layer, podocyte calcium signaling, anti-inflammatory and metabolic functions and enhancing endogenous tissue repair. The greater efficacy of sparsentan compared to ARB on multiple aspects of renal pathophysiology underscores the importance of the interplay of ET-1 and AngII signaling in the pathogenesis and therapy of FSGS.

METHODS

Sex as a biological variable. Our study examined male and female animals, and similar findings are reported for both sexes.

Animals and treatments. Several transgenic mouse models were used in the present study (Fig. 1A) including TRPC6 wild-type (WT, for healthy physiological model) or transgenic mice with podocyte-specific TRPC6 overexpression (Pod-TRPC6 TG, for FSGS model) that were developed and previously characterized (5). Both models were intercrossed with mice expressing either GCaMP5/tdTomato (the intensely green and highly calcium sensitive fluorescent protein GCaMP5G and the calcium insensitive red fluorescent protein tdTomato)(52) or the multicolor Confetti construct (membrane-targeted CFP, nuclear GFP, cytosolic YFP or cytosolic RFP)(53) conditionally in either podocytes (Pod)(54), cells of the renin lineage (Ren1d, which include vascular smooth muscle, JG renin, mesangial and parietal epithelial cells)(55), or endothelial cells (Cdh5)(56) using Cre/lox technology as published recently (18, 24, 25, 30, 45). Some of the presently used transgenic mice were provided by academic investigators, such as the Cdh5(PAC)-CreERT2 mice (by Ralf Adams, University of Münster, Münster, Germany, Cancer Research UK Scientist via Cancer Research Technology Limited), and the Ren1d-Cre mice (Ariel Gomez, University of Virginia, Charlottesville, VA). None of these mice except Pod-TRPC6 TG showed any morphological or functional abnormalities compared to WT mice (not shown). Mouse breeding pairs were purchased from the Jackson Laboratory (Bar Harbor, ME) and were bred to homozygosity for all transgenes and maintained at the University of Southern California in specific pathogen-free quarters according to a homozygous/hemizygous breeding scheme. Tamoxifen was

administered by oral gavage (75 mg/kg body wt) once at 4 weeks of age in Cdh5-Confetti mouse models resulting in endothelial cell-specific expression of reporters (30). Equal number of male and female mice at 6-8 weeks of age (healthy physiological model) or at six months (Ren1d-Confetti and Cdh5-Confetti TRPC6 TG FSGS model) or 1.5 years of age (Pod-GCaMP5/tdTomato TRPC6 TG FSGS model) were used in the present study in three separate groups (n=8 each): received either no drug control, sparsentan (120 mg/kg body wt), or losartan (10 mg/kg body wt) daily in custom rodent chow for two or six weeks of follow-up as indicated. Some animals received ET-1 (50 ng/kg body wt) or Ang II (400 ng/kg body wt) or both injected in bolus into the cannulated carotid artery during intravital imaging or blood pressure measurements (described below). At the end of the treatment period, anesthetized mice were perfusion fixed through the heart and the kidneys were harvested for histological analysis. Animal studies were performed in accordance with guidelines approved by the Institutional Animal Care and Use Committee at the University of Southern California.

Intravital multiphoton microscopy (MPM). Under continuous anesthesia (Isoflurane 1–4% inhalant via nose-cone), the left kidney was exteriorized through a flank incision and the animals were placed on the stage of an inverted microscope with the exposed kidney placed in a coverslip-bottomed chamber bathed in normal saline as described previously (20, 57). Body temperature was maintained with a homeothermic blanket system (Harvard Apparatus). Alexa Fluor 680-conjugated bovine serum albumin (Thermo Fisher, Waltham, MA) was administered iv. by retro-orbital injections to label the circulating plasma (30 μ L iv. bolus from 10 μ g/ml stock solution). The images were acquired using a Leica SP8 DIVE multiphoton confocal fluorescence imaging system with a 40 \times Leica water-immersion objective (numerical aperture 1.1) powered by a

Chameleon Discovery laser at 960 nm (Coherent, Santa Clara, CA) and a DMI8 inverted microscope's external Leica 4Tune spectral hybrid detectors (emission at 510-530 nm for GCaMP5 and at 580-600 nm for tdTomato, and 475-485 nm for detecting second harmonic generation, SHG) (Leica Microsystems, Heidelberg, Germany). When imaging Confetti+ cells, CFP, GFP, YFP and RFP emission was detected at 473, 514, 545 and 585 nm, respectively. The potential toxicity of laser excitation and fluorescence to the cells was minimized by using a low laser power and high scan speeds to keep total laser exposure as minimal as possible. The usual image acquisition (12-bit, 512×512 pixel) consisted of only one z stack per tissue volume (<1 min), which resulted in no apparent cell injury. Fluorescence images were collected in time series (xyt, 526 ms per frame) with the Leica LAS X imaging software and using the same instrument settings (laser power, offset, gain of both detector channels). The strong, positive, cell-specific signal (GCaMP5-tdTomato fluorescence) and high-resolution MPM imaging allowed for easy identification of AA and EA VSMCs, mesangial, and tubular cells. Regions of interest (ROIs) were drawn closely over the total cell body of single cells and the changes in the ratio of mean GCaMP5 and tdTomato F/F₀ fluorescence intensity (ratio was normalized to baseline) were measured in the defined ROI as an index of intracellular calcium changes using the Quantify package of LAS X software (3.6.0.20104; Leica-Microsystems).

Glomerular hemodynamics. Alexa Fluor 680-conjugated bovine serum albumin was used to label the circulating plasma and the negative labeled (albumin-excluding) red blood cells (RBCs). Quantitative imaging of single nephron glomerular filtration rate (SNGFR, by measuring the volume/transit time of a systemically injected Lucifer Yellow dye bolus in the early proximal tubule), glomerular diameter, afferent (AA) and efferent (EA) diameter was performed as

described previously (57, 58). Red blood cell velocity (RBCV) in glomerular capillaries was measured using line (xt) scans of capillary lumen as described before (57). GFB function was evaluated based on measurement of albumin leakage into the Bowman's space (glomerular sieving coefficient, GSC of albumin) (22, 57). ROIs were drawn in glomerular capillary plasma and the Bowman's space, and image analysis was performed as described previously (23, 34).

Blood pressure measurements. Blood pressure was measured by either tail-cuff plethysmography using a Visitech BP-2000 system (Visitech Systems, Apex, NC) or directly with a pressure transducer via the cannulated carotid artery in anesthetized animals using an analog single-channel transducer signal conditioner model BP-1 (World Precision Instruments) as described before (59, 60). All transducer measurement were taken in late afternoon hours at the end of the inactive, resting (non-feeding) phase of the animal's circadian cycle. Tail-cuff measurements were performed in the early morning, and all animals underwent a training period of 5 days before the start of experimental measurements.

Endothelial surface layer (GEC glycocalyx), mitochondrial membrane potential, and immune cell imaging. FITC-labeled wheat germ agglutinin (FITC-WGA) lectin (*Triticum vulgare*; L4895, Sigma, MO), administered via retro-orbital sinus at 2 μ g/g body weight was used to visualize the entire glomerular endothelial glycocalyx. FITC-WGA lectin-positive region of the glomerular endothelial capillary surface was visible immediately after injection. Quantification of glycocalyx thickness was performed on capillary wall line profiles by calculating the width of FITC-WGA signal at half-maximum fluorescence intensity as described before (19, 23). In case of capillary segment heterogeneity, areas with the maximum glycocalyx

thickness were measured. To quantitatively visualize mitochondrial membrane potential and glomerular immune cell homing, MitoTracker Red (ThermoFisher Scientific, Catalog number M7512 dissolved in DMSO) and anti-mouse/human CD44-Alexa Fluor 488 antibodies (BioLegend, Catalog number 103016) were injected iv via retro-orbital sinus, respectively, and imaged as described before (21, 31, 61).

Physiologic and biochemical measurements. Spot urine was collected from animals, and urine albumin was measured by using murine microalbuminuria ELISA kit (Albuwell M kits, Exocell, Philadelphia, PA, USA). Urine creatinine was measured via microplate assay (The Creatinine Companion, Exocell, Philadelphia, PA, USA), and albumin/creatinine ratio (ACR) was calculated.

Tissue processing, immunohistochemistry and histology. After anesthesia with a combination of ketamine (100 mg per kg body weight) and xylazine (10 mg per kg body weight), animals were perfused with ice cold PBS into the left ventricle followed by ice cold 4% PFA for 2 minutes each, and tissues were fixed by 4% PFA at 4°C overnight. To visualize Confetti colors, tissues were embedded in OCT after sucrose cryoprotection method (30% sucrose at room temperature for 3 hours) and flash frozen. Cryosections (18 µm thickness) were imaged using the same Leica TCS SP8 microscope as above. Confetti⁺ clonal or unicolor tracing units were defined as numerous directly adjacent individual cells that featured the same Confetti color combination. All ten possible Confetti color combinations were observed as described before (30). The counting of Confetti⁺ cells and clones was facilitated by standardized image thresholding using ImageJ (NIH), Leica LAS X (Leica Microsystems Inc.), and cell-counting algorithms of Imaris 9.2 3D

image visualization and analysis software (Bitplane USA) for imaging same-size Z-stacks. Immunofluorescence (IF) staining was performed on paraffin sections (6 μm thickness). After antigen retrieval (8 min at 95°C in citrate buffer using pressure cooker) and blocking (30 min in goat blocking buffer), the sections were incubated with anti-p57 (1:100, ab228635, Abcam, Cambridge MA, USA) or WT1 (1:100, ab89901, Abcam) primary antibodies followed by incubation with the secondary antibodies conjugated with Alexa 594 (1:500, A-11012, Invitrogen, CA, USA). Slides were mounted by using DAPI-containing mounting media (VectaShield, Vector Laboratories Inc., Burlingame, CA). Some paraffin sections were stained using Picrosirius Red and imaged using the same Leica TCS SP8 microscope as above. Images were analyzed using Image J software (NIH) for measuring Picrosirius Red pixel density per unit area as an index of glomerulosclerosis and tubulointerstitial fibrosis.

Statistics. Data represent average \pm SEM and were analyzed using one-way ANOVA for multiple groups with post-hoc comparison by Tukey's or Sidak's tests as appropriate. $P < 0.05$ was considered significant. Statistical analyses were performed using GraphPad Prism 9.0c (GraphPad Software, Inc.).

Study approval. All animal protocols were approved by the Institutional Animal Care and Use Committee at the University of Southern California (Los Angeles, CA) and followed the NIH Guide for the Care and Use of Laboratory Animals.

Data availability. All supporting values for this manuscript are provided in the "Supporting Data Values" XLS file.

AUTHOR CONTRIBUTIONS

GG, UNS, SD, AI, and JPP conducted the experiments; GG, RK, PWB, and JPP designed the study and wrote the paper. All authors approved the final version of the manuscript. Co-author employees of Travers Therapeutics were not involved in data analysis.

ACKNOWLEDGMENTS

This work was supported by a research collaboration grant by Travers Therapeutics and by NIH S10 OD021833 to the USC Multi-Photon Microscopy Core.

REFERENCES

1. Saran R, Robinson B, Abbott KC, Bragg-Gresham J, Chen X, Gipson D, et al. US Renal Data System 2019 Annual Data Report: Epidemiology of Kidney Disease in the United States. *Am J Kidney Dis.* 2020;75(1 Suppl 1):A6-a7.
2. Campbell KN, and Tumlin JA. Protecting Podocytes: A Key Target for Therapy of Focal Segmental Glomerulosclerosis. *Am J Nephrol.* 2018;47 Suppl 1(Suppl 1):14-29.
3. Feng D, DuMontier C, and Pollak MR. Mechanical challenges and cytoskeletal impairments in focal segmental glomerulosclerosis. *Am J Physiol Renal Physiol.* 2018;314(5):F921-925.
4. Dryer SE, and Reiser J. TRPC6 channels and their binding partners in podocytes: role in glomerular filtration and pathophysiology. *Am J Physiol Renal Physiol.* 2010;299(4):F689-701.
5. Krall P, Canales CP, Kairath P, Carmona-Mora P, Molina J, Carpio JD, et al. Podocyte-specific overexpression of wild type or mutant *trpc6* in mice is sufficient to cause glomerular disease. *PLoS One.* 2010;5(9):e12859.
6. Reiser J, Polu KR, Möller CC, Kenlan P, Altintas MM, Wei C, et al. TRPC6 is a glomerular slit diaphragm-associated channel required for normal renal function. *Nat Genet.* 2005;37(7):739-44.
7. Winn MP, Conlon PJ, Lynn KL, Farrington MK, Creazzo T, Hawkins AF, et al. A mutation in the TRPC6 cation channel causes familial focal segmental glomerulosclerosis. *Science.* 2005;308(5729):1801-4.
8. Eckel J, Lavin PJ, Finch EA, Mukerji N, Burch J, Gbadegesin R, et al. TRPC6 enhances angiotensin II-induced albuminuria. *J Am Soc Nephrol.* 2011;22(3):526-35.

9. Komers R, and Plotkin H. Dual inhibition of renin-angiotensin-aldosterone system and endothelin-1 in treatment of chronic kidney disease. *Am J Physiol Regul Integr Comp Physiol*. 2016;310(10):R877-84.
10. Kowala MC, Murugesan N, Tellew J, Carlson K, Monshizadegan H, Ryan C, et al. Novel dual action AT1 and ETA receptor antagonists reduce blood pressure in experimental hypertension. *J Pharmacol Exp Ther*. 2004;309(1):275-84.
11. Trachtman H, Nelson P, Adler S, Campbell KN, Chaudhuri A, Derebail VK, et al. DUET: A Phase 2 Study Evaluating the Efficacy and Safety of Sparsentan in Patients with FSGS. *J Am Soc Nephrol*. 2018;29(11):2745-54.
12. Campbell KN, Gesualdo L, Murphy E, Rheault MN, Srivastava T, Tesar V, et al. Sparsentan for Focal Segmental Glomerulosclerosis in the DUET Open-Label Extension: Long-term Efficacy and Safety. *Kidney Med*. 2024;6(6):100833.
13. Kohan DE, Bedard PW, Jenkinson C, Hendry B, and Komers R. Mechanism of protective actions of sparsentan in the kidney: lessons from studies in models of chronic kidney disease. *Clin Sci (Lond)*. 2024;138(11):645-62.
14. Rheault MN, Alpers CE, Barratt J, Bieler S, Canetta P, Chae DW, et al. Sparsentan versus Irbesartan in Focal Segmental Glomerulosclerosis. *N Engl J Med*. 2023;389(26):2436-45.
15. Rovin BH, Barratt J, Heerspink HJL, Alpers CE, Bieler S, Chae DW, et al. Efficacy and safety of sparsentan versus irbesartan in patients with IgA nephropathy (PROTECT): 2-year results from a randomised, active-controlled, phase 3 trial. *Lancet*. 2023;402(10417):2077-90.
16. Reily C, Moldoveanu Z, Pramparo T, Hall S, Huang ZQ, Rice T, et al. Sparsentan ameliorates glomerular hypercellularity and inflammatory-gene networks induced by IgA1-IgG immune complexes in a mouse model of IgA nephropathy. *Am J Physiol Renal Physiol*. 2024;326(5):F862-875.
17. Burford JL, Gyarmati G, Shirato I, Kriz W, Lemley KV, and Peti-Peterdi J. Combined use of electron microscopy and intravital imaging captures morphological and functional features of podocyte detachment. *Pflugers Arch*. 2017;469(7-8):965-74.
18. Burford JL, Villanueva K, Lam L, Riquier-Brison A, Hackl MJ, Pippin J, et al. Intravital imaging of podocyte calcium in glomerular injury and disease. *J Clin Invest*. 2014;124(5):2050-8.
19. Butler MJ, Ramnath R, Kadoya H, Desposito D, Riquier-Brison A, Ferguson JK, et al. Aldosterone induces albuminuria via matrix metalloproteinase-dependent damage of the endothelial glycocalyx. *Kidney Int*. 2019;95(1):94-107.
20. Hackl MJ, Burford JL, Villanueva K, Lam L, Susztak K, Schermer B, et al. Tracking the fate of glomerular epithelial cells in vivo using serial multiphoton imaging in new mouse models with fluorescent lineage tags. *Nat Med*. 2013;19(12):1661-6.
21. Kadoya H, Yu N, Schiessl IM, Riquier-Brison A, Gyarmati G, Desposito D, et al. Essential role and therapeutic targeting of the glomerular endothelial glycocalyx in lupus nephritis. *JCI Insight*. 2020;5(19):e131252.
22. Peti-Peterdi J, and Sipos A. A high-powered view of the filtration barrier. *J Am Soc Nephrol*. 2010;21(11):1835-41.
23. Salmon AH, Ferguson JK, Burford JL, Gevorgyan H, Nakano D, Harper SJ, et al. Loss of the endothelial glycocalyx links albuminuria and vascular dysfunction. *J Am Soc Nephrol*. 2012;23(8):1339-50.

24. Kaverina NV, Kadoya H, Eng DG, Rusiniak ME, Sequeira-Lopez ML, Gomez RA, et al. Tracking the stochastic fate of cells of the renin lineage after podocyte depletion using multicolor reporters and intravital imaging. *PLoS One*. 2017;12(3):e0173891.
25. Schiessl IM, Fremter K, Burford JL, Castrop H, and Peti-Peterdi J. Long-Term Cell Fate Tracking of Individual Renal Cells Using Serial Intravital Microscopy. *Methods Mol Biol*. 2020;2150:25-44.
26. Shroff UN, Schiessl IM, Gyarmati G, Riquier-Brison A, and Peti-Peterdi J. Novel fluorescence techniques to quantitate renal cell biology. *Methods Cell Biol*. 2019;154:85-107.
27. Peti-Peterdi J, Kidokoro K, and Riquier-Brison A. Novel in vivo techniques to visualize kidney anatomy and function. *Kidney Int*. 2015;88(1):44-51.
28. Gyarmati G, Shroff UN, Riquier-Brison A, Desposito D, Ju W, Stocker SD, et al. Neuronally differentiated macula densa cells regulate tissue remodeling and regeneration in the kidney. *J Clin Invest*. 2024;134(11):e174558.
29. Kidokoro K, Cherney DZI, Bozovic A, Nagasu H, Satoh M, Kanda E, et al. Evaluation of Glomerular Hemodynamic Function by Empagliflozin in Diabetic Mice Using In Vivo Imaging. *Circulation*. 2019;140(4):303-15.
30. Desposito D, Schiessl IM, Gyarmati G, Riquier-Brison A, Izuhara AK, Kadoya H, et al. Serial intravital imaging captures dynamic and functional endothelial remodeling with single-cell resolution. *JCI Insight*. 2021;6(10):e123392.
31. Gyarmati G, Shroff UN, Izuhara A, Hou X, Da Sacco S, Sedrakyan S, et al. Intravital imaging reveals glomerular capillary distension and endothelial and immune cell activation early in Alport syndrome. *JCI Insight*. 2022;7(1):e152676.
32. Kitching AR, and Hickey MJ. Immune cell behaviour and dynamics in the kidney - insights from in vivo imaging. *Nat Rev Nephrol*. 2022;18(1):22-37.
33. Molitoris BA, Dunn KW, and Sandoval RM. Proximal tubule role in albumin homeostasis: controversy, species differences, and the contributions of intravital microscopy. *Kidney Int*. 2023;104(6):1065-9.
34. Nakano D, Kobori H, Burford JL, Gevorgyan H, Seidel S, Hitomi H, et al. Multiphoton imaging of the glomerular permeability of angiotensinogen. *J Am Soc Nephrol*. 2012;23(11):1847-56.
35. Nagasawa H, Suzuki H, Jenkinson C, Ueda S, Fukao Y, Nakayama M, et al. MO261: Sparsentan, The Dual Endothelin Angiotensin Receptor Antagonist (DEARA), Attenuates Albuminuria and Protects from the Development of Renal Injury to a Greater Extent Than Losartan in the GDDY Mouse Model of IGA Nephropathy: A 16-Week Study. *Nephrology Dialysis Transplantation*. 2022;37(Supplement_3).
36. Reily C, Moldoveanu Z, Pramparo T, Hall SD, Novak L, Komers R, et al. The Dual Endothelin Angiotensin Receptor Antagonist (DEARA) Sparsentan Protects from Glomerular Hypercellularity and Associated Immune/Inflammatory Gene Network Activity in a Model of IgA Nephropathy [Abstract]. *J Am Soc Nephrol*. 2021;32:PO1454.
37. Anguiano L, Riera M, Pascual J, and Soler MJ. Endothelin Blockade in Diabetic Kidney Disease. *J Clin Med*. 2015;4(6):1171-92.
38. Harrison-Bernard LM, Navar LG, Ho MM, Vinson GP, and el-Dahr SS. Immunohistochemical localization of ANG II AT1 receptor in adult rat kidney using a monoclonal antibody. *Am J Physiol*. 1997;273(1 Pt 2):F170-7.

39. Lichtnekert J, Kaverina NV, Eng DG, Gross KW, Kutz JN, Pippin JW, et al. Renin-Angiotensin-Aldosterone System Inhibition Increases Podocyte Derivation from Cells of Renin Lineage. *J Am Soc Nephrol*. 2016;27(12):3611-27.
40. Pippin JW, Kaverina NV, Eng DG, Krofft RD, Glenn ST, Duffield JS, et al. Cells of renin lineage are adult pluripotent progenitors in experimental glomerular disease. *Am J Physiol Renal Physiol*. 2015;309(4):F341-58.
41. Lasagni L, Angelotti ML, Ronconi E, Lombardi D, Nardi S, Peired A, et al. Podocyte Regeneration Driven by Renal Progenitors Determines Glomerular Disease Remission and Can Be Pharmacologically Enhanced. *Stem Cell Reports*. 2015;5(2):248-63.
42. Shankland SJ, Anders HJ, and Romagnani P. Glomerular parietal epithelial cells in kidney physiology, pathology, and repair. *Curr Opin Nephrol Hypertens*. 2013;22(3):302-9.
43. Kaverina NV, Eng DG, Largent AD, Daehn I, Chang A, Gross KW, et al. WT1 Is Necessary for the Proliferation and Migration of Cells of Renin Lineage Following Kidney Podocyte Depletion. *Stem Cell Reports*. 2017;9(4):1152-66.
44. McMahon EJ, Bauer JD, Hawley CM, Isbel NM, Stowasser M, Johnson DW, et al. A randomized trial of dietary sodium restriction in CKD. *J Am Soc Nephrol*. 2013;24(12):2096-103.
45. Manavski Y, Lucas T, Glaser SF, Dorsheimer L, Günther S, Braun T, et al. Clonal Expansion of Endothelial Cells Contributes to Ischemia-Induced Neovascularization. *Circ Res*. 2018;122(5):670-7.
46. Anderson M, Roshanravan H, Khine J, and Dryer SE. Angiotensin II activation of TRPC6 channels in rat podocytes requires generation of reactive oxygen species. *J Cell Physiol*. 2014;229(4):434-42.
47. Ilatovskaya DV, Blass G, Palygin O, Levchenko V, Pavlov TS, Grzybowski MN, et al. A NOX4/TRPC6 Pathway in Podocyte Calcium Regulation and Renal Damage in Diabetic Kidney Disease. *J Am Soc Nephrol*. 2018;29(7):1917-27.
48. Daehn I, Casalena G, Zhang T, Shi S, Fenninger F, Barasch N, et al. Endothelial mitochondrial oxidative stress determines podocyte depletion in segmental glomerulosclerosis. *J Clin Invest*. 2014;124(4):1608-21.
49. Barton M, and Sorokin A. Endothelin and the glomerulus in chronic kidney disease. *Semin Nephrol*. 2015;35(2):156-67.
50. Ebefors K, Wiener RJ, Yu L, Azeloglu EU, Yi Z, Jia F, et al. Endothelin receptor-A mediates degradation of the glomerular endothelial surface layer via pathologic crosstalk between activated podocytes and glomerular endothelial cells. *Kidney Int*. 2019;96(4):957-70.
51. Boels MG, Avramut MC, Koudijs A, Dane MJ, Lee DH, van der Vlag J, et al. Atrasentan Reduces Albuminuria by Restoring the Glomerular Endothelial Glycocalyx Barrier in Diabetic Nephropathy. *Diabetes*. 2016;65(8):2429-39.
52. Gee JM, Smith NA, Fernandez FR, Economo MN, Brunert D, Rothermel M, et al. Imaging activity in neurons and glia with a Polr2a-based and cre-dependent GCaMP5G-IRES-tdTomato reporter mouse. *Neuron*. 2014;83(5):1058-72.
53. Snippert HJ, van der Flier LG, Sato T, van Es JH, van den Born M, Kroon-Veenboer C, et al. Intestinal crypt homeostasis results from neutral competition between symmetrically dividing Lgr5 stem cells. *Cell*. 2010;143(1):134-44.
54. Moeller MJ, Sanden SK, Soofi A, Wiggins RC, and Holzman LB. Podocyte-specific expression of cre recombinase in transgenic mice. *Genesis*. 2003;35(1):39-42.

55. Sequeira López ML, Pentz ES, Nomasa T, Smithies O, and Gomez RA. Renin cells are precursors for multiple cell types that switch to the renin phenotype when homeostasis is threatened. *Dev Cell*. 2004;6(5):719-28.
56. Wang Y, Nakayama M, Pitulescu ME, Schmidt TS, Bochenek ML, Sakakibara A, et al. Ephrin-B2 controls VEGF-induced angiogenesis and lymphangiogenesis. *Nature*. 2010;465(7297):483-6.
57. Kang JJ, Toma I, Sipos A, McCulloch F, and Peti-Peterdi J. Quantitative imaging of basic functions in renal (patho)physiology. *Am J Physiol Renal Physiol*. 2006;291(2):F495-502.
58. Peti-Peterdi J, Kidokoro K, and Riquier-Brison A. Intravital imaging in the kidney. *Current opinion in nephrology and hypertension*. 2016;25(3):168-73.
59. Riquier-Brison ADM, Sipos A, Prókai Á, Vargas SL, Toma L, Meer EJ, et al. The macula densa prorenin receptor is essential in renin release and blood pressure control. *Am J Physiol Renal Physiol*. 2018;315(3):F521-534.
60. Pluznick JL, Protzko RJ, Gevorgyan H, Peterlin Z, Sipos A, Han J, et al. Olfactory receptor responding to gut microbiota-derived signals plays a role in renin secretion and blood pressure regulation. *Proc Natl Acad Sci U S A*. 2013;110(11):4410-5.
61. Gyarmati G, Kadoya H, Moon JY, Burford JL, Ahmadi N, Gill IS, et al. Advances in Renal Cell Imaging. *Semin Nephrol*. 2018;38(1):52-62.

FIGURES AND FIGURE LEGENDS




Figure 1. General study design and the effects of sparsentan and losartan on glomerular hemodynamics in the control healthy Ren1d-GCaMP5/tTomato mouse kidney. (A) Schematic illustration of the specific transgenic mouse strains used for either physiological (6-8 weeks old) and FSGS disease models (6 months – 1.5 years old)(left panel), duration of treatment and disease development (center panel) and the applied MPM imaging and classic phenotyping readouts (right panel) in each model. **(B)** Intravital MPM images of control healthy Ren1d-GCaMP5/tTomato mouse kidney glomeruli (G) with their afferent (AA) and efferent (EA) arterioles from no drug control, sparsentan, and losartan treated mice as indicated. The circulating plasma was labeled by iv injected Albumin-Alexa Fluor 680 (grey). Note the presence of the genetically encoded calcium reporter GCaMP5 (green) and the calcium-insensitive tdTomato (red) in cells of the renin lineage including AA and EA VSMCs, and extraglomerular and

intraglomerular mesangial cells (arrowheads). **(C-I)** Statistical summary of the measured hemodynamic parameters (n=8 each) including AA (**C**), EA (**D**), and glomerular diameters (**E**), glomerular tuft area (**F**), glomerular capillary diameter (**G**), single nephron glomerular filtration rate (SNGFR, **H**) and glomerular capillary red blood cell velocity (RBCV, **I**). Data represent mean \pm SEM, *P< 0.05, **P<0.01, ***P<0.001, ****P< 0.0001 using one-way ANOVA with Tukey's multiple comparison test, for all panels n=10 measurements averaged for each of the n=8 mice (n=4 males (blue) and n=4 females (red)) in each group. Bar is 20 μ m for all panels.




Figure 2. Effects of sparsentan and losartan on glomerular hemodynamic alterations caused by acute agonist-induced vasoconstriction in control healthy Ren1d-GCaMP5/tdTomato mice. (A) Intravital MPM images of glomeruli (G) with their afferent (AA) and efferent (EA) arterioles before (top panel) and after (bottom panel) the bolus injection of endothelin (ET-1, 50 ng/kg) with AngII (400 ng/kg) combined into the cannulated carotid artery in no drug control (left), sparsentan (center), and losartan (right) treated mice. The circulating plasma was labeled by iv injected Albumin-Alexa Fluor 680 (grey). Note the presence of the genetically encoded calcium reporter GCaMP5 (green) and the calcium-insensitive tdTomato (red) in cells of the renin lineage including AA and EA VSMCs, and extraglomerular and intraglomerular mesangial cells (arrowheads). **(B)** Time-lapse recordings of AA VSMC calcium (GCaMP5/tdTomato fluorescence ratio) changes in response to bolus ET-1+AngII injection into the carotid artery

(vertical line) in the three experimental groups. **(C-F)** Statistical summary of the measured hemodynamic parameters (n=8 each) normalized to baseline (ratio of maximum effect/before injection), including GCaMP5/tdTomato fluorescence ratio (F_{\max}/F_0) in VSMCs of the AA **(C)**, and the AA **(D)**, and glomerular diameters **(E)**, and glomerular tuft area **(F)**. Data represent mean \pm SEM, *P< 0.05, **P<0.01, ***P<0.001, ****P< 0.0001 using one-way ANOVA with Sidak's multiple comparisons test, for all panels n=10 measurements averaged for each of the n=8 mice (n=4 males (blue) and n=4 females (red)) in each group. Bar is 20 μ m for all panels.




Figure 3. Effects of sparsentan and losartan on glomerular and vascular tissue remodeling in control healthy conditions. (A, F) Ren1d-Confetti (A) and Cdh5-Confetti (F) mouse kidney sections from no drug control (left), sparsentan (center), or losartan (right) treatment. Note the presence of the genetically encoded multi-color Confetti reporter (membrane CFP (blue), nuclear GFP (green), cytosolic YFP (yellow) and RFP (red)) in cells of the renin lineage (including afferent arteriole (AA) vascular smooth muscle and renin cells, extraglomerular and intraglomerular mesangial cells (arrowheads), parietal epithelial cells and podocytes of the

Bowman's capsule (arrows) in A) and in glomerular (G), AA, and efferent arteriole (EA) vascular endothelium (F). **(B-J)** Statistical summary (n=8 each) of the number of Ren1d and Cdh5-Confetti+ cells **(B, G)**, identical color cell groups (clones, blue/yellow arrows in F) **(C, H)**, cells per clone **(D, I)** and clone frequency **(E, J)** in the various treatment groups. **(K)** Confetti+ cortical (top) and medullary (bottom) tubular cells in Ren1d-Confetti mouse kidney sections from no drug control (left), sparsentan (center), or losartan (right) treatment. Note the presence or absence of clonal (identical color) cell groups in the proximal tubule (PT, inset), cortical (CCD) and medullary collecting duct (CD). Bars are 20 μ m. **(L)** Statistical summary (n=8 each) of clone frequency in renal tubules in Ren1d-Confetti mouse kidney sections in the various treatment groups. Data represent mean \pm SEM, *P< 0.05, **P<0.01, ***P<0.001, ****P< 0.0001, ns: not significant, using one-way ANOVA with Tukey's multiple comparison test, for all panels n=10 measurements averaged for each of the n=8 mice (n=4 males and n=4 females) in each group.




Figure 4. Effects of sparsentan and losartan on glomerular hemodynamics and GFB function in FSGS. (A) Intravital MPM images of glomeruli (G) and their afferent (AA) and efferent (EA) arterioles from no drug control (left), sparsentan (center), and losartan (right) treated Pod-GCaMP5/tdTomato TRPC6 TG mice (1.5 years old). The circulate ng plasma was labeled by iv injected Albumin-Alexa Fluor 680 (grey). Note the presence of the genetically encoded calcium reporter GCaMP5 (green) and the calcium-insensitive tdTomato (red) in podocytes. Podocytes with high calcium (intense green) are visible in a segmental pattern (arrows) adjacent to areas of adhesions between capillaries and parietal podocytes (arrowheads). Plasma albumin leaked through the GFB is visible in the Bowman's space (asterisk). (B-J) Statistical summary of the measured hemodynamic and GFB parameters in the various treatment groups (n=8 each)

including podocyte calcium (based on the ratio of GCaMP5/tdTomato fluorescence (F) intensity, **B**), AA (**C**), EA (**D**), and glomerular diameters (**E**), glomerular tuft area (**F**), single nephron glomerular filtration rate (SNGFR, **G**), glomerular capillary red blood cell velocity (RBCV, **H**), albumin leakage through the GFB (based on albumin glomerular sieving coefficient (GSC), **I**), and albuminuria (based on urinary albumin/creatinine ratio (ACR) normalized to baseline, **J**). (**K-L**) Systolic blood pressure (BP) at baseline (**K**) and elevations (Δ BP) in response to acute AngII injection (**L**)(n=8 each). Data represent mean \pm SEM, *P< 0.05, **P<0.01, ***P<0.001, ****P< 0.0001, ns=not significant, using one-way ANOVA with Tukey's multiple comparison test, for all panels n=10 measurements averaged for image-based parameters (panels B-I) for each of the n=8 mice (n=4 males (blue) and n=4 females (red)) in each group. Bar is 20 μ m for all panels.




Figure 5. Effects of sparsentan and losartan on alterations in glomerular hemodynamics and GFB function caused by acute agonist-induced vasoconstriction in FSGS. (A) Intravital MPM images of glomeruli (G) with their afferent (AA) and efferent (EA) arterioles before (top) and after (bottom) the bolus injection of ET-1 (50 ng/kg) and AngII (400 ng/kg) combined into the cannulated carotid artery in no drug control (left), sparsentan (center), and losartan (right) treated (for 6 weeks) Pod-GCaMP5/tdTomato TRPC6 TG mice (1.5 years old). The circulating plasma was labeled by iv injected Albumin-Alexa Fluor 680 (grey), which also illuminated the AA and EA. Note the presence of the genetically encoded calcium reporter GCaMP5 (green) and the calcium-insensitive tdTomato (red) in podocytes. Insets show the GCaMP5 (green) channel separately to better visualize podocyte calcium changes. (B-E) Statistical summary of the measured hemodynamic and GFB parameters (n=8 each) normalized to baseline (ratio of maximum effect/before injection), including GCaMP5/tdTomato fluorescence ratio (F_{max}/F_0) in podocytes (B), and the AA (C) and glomerular diameters (D), and glomerular tuft area (E). Data represent mean \pm SEM, *P< 0.05, **P<0.01, ***P<0.001, ****P< 0.0001 using one-way ANOVA with Tukey's multiple comparison test, for all panels n=10 measurements averaged for each of the n=8 mice (n=4 males (blue) and n=4 females (red)) in each group. Bars are 20 μ m.




Figure 6. Effects of sparsentan and losartan on podocyte number, glomerulosclerosis and tissue fibrosis, and endogenous tissue regeneration in FSGS. (A-B) p57 (A) or WT1 (B) immunolabeling (red, arrowheads) and statistical summary of p57+ or WT1+ podocyte number/glomerulus in fixed histological sections from no drug control (left), sparsentan (center), and losartan (right) treated (for 6 weeks) Pod-GCaMP5/tdTomato TRPC6 TG mouse (1.5 years old) kidneys (n=8 each). Tissue autofluorescence is shown in yellow. (C) Picrosirius red (red)

staining of kidney sections from the same groups/mice as in A, and glomerulosclerosis index (based on picosirius red density per glomeruli) and tissue fibrosis index (based on picosirius red density per full image frame) in the various treatment groups (n=8 each). **(D-G)** Clonal analysis of Ren1d-Confetti **(D)** and Cdh5-Confetti **(E)** Pod-TRPC6 TG mouse (6 months old) kidneys from same treatment groups as shown in A (n=8 each). Note the presence of the genetically encoded multi-color Confetti reporter (CFP (blue), GFP (green), YFP (yellow) and RFP (red)) in cells of the renin lineage **(D)** or vascular endothelium **(E)** and entirely clonal (identical color) glomeruli (yellow/magenta/blue arrows) in sparsentan **(C, center)** and to a lower extent in losartan treated mice **(C, right)**. Statistical summary of the number of Confetti+ cells, identical color cell groups, cells per clone and clone frequency **(F, G)** in the various treatment groups in Ren1d **(D, F)** and Cdh5-Confetti mice **(E, G)**. Data represent mean \pm SEM, *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001, ns=not significant, using one-way ANOVA with Tukey's multiple comparison test, for all panels n=10 measurements averaged for each of the n=8 mice (n=4 males (blue) and n=4 females (red)) in each group. G: glomeruli. Bars are 20 μ m.




Figure 7. Effects of sparsentan and losartan on glomerular cell metabolism, endothelial surface layer, and immune cell homing in FSGS. (A-B) Intravital MPM images and quantitative analysis of MitoTracker Red (A, magenta) and FITC-WGA (B, green) fluorescence in glomeruli (G) from no drug control (left), sparsentan (center), and losartan (right) treated (for 6 weeks) Pod-TRPC6 TG mouse (6 months old) kidneys. Note the intensely labeled podocytes suggesting high level of oxidative stress (A, arrowheads) and endothelial glycocalyx heterogeneity (B, accumulation in some capillary segments (thick arrows) but undetectable levels in other regions (thin arrows) in control but not in sparsentan treated mice. Numerous CD44⁺ (green) immune cells (B, arrowheads) were found in the lumen of glomerular capillaries in control FSGS mice. Statistical summary of MitoTracker Red intensity in the entire glomerulus or selectively in podocytes (A) and glomerular endothelial surface layer (glycocalyx) thickness and CD44⁺ immune cell homing (B) in the various treatment groups (n=8 each). Data represent mean \pm SEM, **P<0.01, ***P<0.001, ****P<0.0001, ns=not significant, using one-way ANOVA with Tukey's multiple comparison test, for all panels n=10 measurements averaged for each of the n=8 mice (n=4 males (blue) and n=4 females (red)) in each group. Bars are 20 μ m.