

SUPPLEMENTAL TABLE


Table S1. Least squares mean change from baseline in primary and secondary efficacy outcome measures on day 8 and study end (visit 4, day 15), carbetocin versus placebo

Outcome measure	Day 8		Day 15	
	LS Mean Change	P Value	LS Mean Change	P Value
HPWSQ-R total score	−5.2	0.0632	−6.7	0.0290
Behavior	−1.9	0.1153	−2.0	0.1172
Drive	−2.1	0.0625	−1.6	0.1436
Severity	−1.1	0.0572	−1.5	0.0248
HPWSQ-R-C total score	−10.2	0.0017	−10.5	0.0014
Behavior	−5.4	0.0017	−4.3	0.0068
Drive	−3.4	0.0019	−2.6	0.0340
Severity	−1.7	0.0061	−2.2	0.0014
CGI-I	−0.9	0.0162	−0.8	0.0233
CY-BOCS	−2.7	0.0956	−6.2	0.0047
Food domain score of Reiss Profile	−3.0	0.0184	−4.4	0.0132

CGI-I, Clinical Global Impressions–Improvement; CY- BOCS, Children’s Yale-Brown Obsessive Compulsive Scale; HPWSQ-R, parent/caregiver-rated Hyperphagia in Prader-Willi Syndrome Questionnaire-Responsiveness; HPWSQ-R-C, clinician-rated Hyperphagia in Prader-Willi Syndrome Questionnaire-Responsiveness; LS, least squares.


SUPPLEMENTAL FIGURES

Figure S1. Change from baseline in HPWSQ-C total score at day 8 and study end (visit 4, day 15). The analysis included all participants with an HPWSQ-C assessment at baseline, day 8, and/or day 15 and was performed using an ANCOVA model with treatment group and study site as fixed effects.




ANCOVA, analysis of covariance; HPWSQ-C, clinician-rated Hyperphagia in Prader-Willi Syndrome Questionnaire-Responsiveness; LS, least squares; SE, standard error.

Figure S2. Change from baseline in CY-BOCS total score at day 8 and study end (visit 4, day 15). The analysis included all participants with a CY-BOCS assessment at baseline, day 8, and/or day 15 and was performed using an ANCOVA model with treatment group and study site as fixed effects.




ANCOVA, analysis of covariance; CY-BOCS, Children's Yale-Brown Obsessive Compulsive Scale; LS, least squares; SE, standard error.

Figure S3. Change from baseline in CGI-I total score at day 8 and study end (visit 4, day 15). The analysis included all participants with a CGI-I assessment at baseline, day 8, and/or day 15 and was performed using an ANCOVA model with treatment group and study site as fixed effects.



ANCOVA, analysis of covariance; CGI-I, Clinical Global Impression–Improvement; LS, least squares; SE, standard error.

Figure S4. Change from baseline in Food domain score of Reiss Profile total score at day 8 and study end (visit 4, day 15). The analysis included all participants with a food domain score of the Reiss Profile assessment at baseline, day 8, and/or day 15 and was performed using an ANCOVA model with treatment group and study site as fixed effects.



ANCOVA, analysis of covariance; LS, least squares; SE, standard error.